

Bianchi Type III String Cosmological Models with Time Dependent Bulk Viscosity

BALI Raj¹ and PRADHAN Anirudh²

¹Department of Mathematics, University of Rajasthan, Jaipur-302 004, India
E-mail : balir5@yahoo.co.in

²Department of Mathematics, Hindu Post-graduate College, Zamania-232 331,
Ghazipur, India
E-mail : pradhan@iucaa.ernet.in, acpradhan@yahoo.com

Abstract

Bianchi type III string cosmological models with bulk viscous fluid for massive string are investigated. To get the determinate model of the universe, we have assumed that the coefficient of bulk viscosity ξ is inversely proportional to the expansion θ in the model and expansion θ in the model is proportional to the shear σ . This leads to $B = \ell C^n$, where ℓ and n are constants. The behaviour of the model in presence and absence of bulk viscosity, is discussed. The physical implications of the models are also discussed in detail.

Keywords: Cosmic string, viscous models, Bianchi type III model
PACS: 98.80.Cq, 04.20.-q

1 Introduction

In recent years, there has been considerable interest in string cosmology. Cosmic strings are topologically stable objects which might be found during a phase transition in the early universe (Kibble [1]). Cosmic strings play an important role in the study of the early universe. These arise during the phase transition after the big bang explosion as the temperature goes down below some critical temperature as predicted by grand unified theories (Zel'dovich et al. [2]; Kibble [1, 3]; Everett [4]; Vilenkin [5]). It is believed that cosmic strings give rise to density perturbations which lead to the formation of galaxies (Zel'dovich [6]). These cosmic strings have stress-energy and couple to the gravitational field. Therefore it is interesting to study the gravitational effects that arise from strings.

The general relativistic treatment of strings was initiated by Letelier [7, 8] and Stachel [9]. Letelier [7] has obtained the solution to Einstein's field equations for a cloud of strings with spherical, plane and cylindrical symmetry. Then, in 1983, he solved Einstein's field equations for a cloud of massive strings and obtained cosmological models in Bianchi I and Kantowski-Sachs space-times. Benerjee et al. [10] have investigated an axially symmetric Bianchi type I string dust cosmological model in presence and absence of magnetic field. The string cosmological models with a magnetic field are also discussed by Chakraborty [11], Tikekar and Patel [12]. Patel and Maharaj [13] investigated stationary rotating world model with magnetic field. Ram and Singh [14] obtained some new exact solutions of string cosmology with and without a source free magnetic field for Bianchi type I space-time in the different basic form considered by Carminati and McIntosh [15]. Exact solutions of string cosmology for Bianchi type II, VI_0 , VIII and IX space-times have been studied by Krori et al. [16] and Wang [17]. Singh and Singh [18] investigated string cosmological models with magnetic field in the context of space-time with G_3 symmetry. Lidsey, Wands and Copeland [19] have reviewed aspects of super string cosmology with the emphasis on the cosmological implications of duality symmetries in the theory. Baysal et al. [20] have investigated the behaviour of a string in the cylindrically symmetric inhomogeneous universe. Bali et al. [21 - 25] have obtained Bianchi types IX, type-V and type-I string cosmological models in general relativity. Yavuz [26] have examined charged strange quark matter attached to the string cloud in the spherical symmetric space-time admitting one-parameter group of conformal motion. Recently Kaluza-Klein cosmological solutions are obtained by Yilmaz [27] for quark matter coupled to the string cloud in the context of general relativity.

On the other hand, the matter distribution is satisfactorily described by perfect fluids due to the large scale distribution of galaxies in our universe. However, a realistic treatment of the problem requires the consideration of material distribution other than the perfect fluid. It is well known that when neutrino decoupling occurs, the matter behaves as a viscous fluid in an early stage of the universe. Viscous fluid cosmological models of early universe have been widely discussed in the literature.

Recently, Seljak et al. [28] have studied cosmological parameters from combining Lyman- α forest with C.M.B. They established the cosmic strings limited to $G\mu < 2.3 \times 10^{-7}$ at 95 percent cathodoluminescence (c.l.), and correlated iso-curvature models are also tightly constrained. Observations on C.M.B. constrain the cosmic string which is predicted in terms of cosmic string power spectrum [29]. However, we are not concerned with this new phenomena. Tikekar and Patel [12], following the techniques used by Letelier and Stachel, obtained some exact Bianchi III cosmological solutions of massive strings in presence of magnetic field. Maharaj et al. [30] have generalized the previous solutions obtained by Tikekar and Patel [12] considering Lie point symmetries. Yadav et al. [31] have studied some Bianchi type I viscous fluid string cosmological models with magnetic field. Recently Wang [32 - 35] has also discussed LRS Bianchi type I and Bianchi type III cosmological models for a cloud string with bulk viscos-

ity. Very recently, Yadav, Rai and Pradhan [36] have found the integrability of cosmic string in Bianchi type III space-time in presence of bulk viscous fluid by applying a new technique. Motivated by the situations discussed above, in this study, we established a formalism for studying the new integrability of massive strings in Bianchi III space-time in presence of a variable bulk viscosity.

2 The Metric and Field Equations

We consider the space-time of general Bianchi III type with the metric

$$ds^2 = -dt^2 + A^2(t)dx^2 + B^2(t)e^{-2\alpha x}dy^2 + C^2(t)dz^2, \quad (1)$$

where α is constant. The energy momentum tensor for a cloud of string dust with a bulk viscous fluid of string is given by Letelier [7] and Landau & Lifshitz [37]

$$T_i^j = \rho v_i v^j - \lambda x_i x^j - \xi v_{;l}^l (g_i^j + v_i v^j), \quad (2)$$

where v_i and x_i satisfy condition

$$v^i v_i = -x^i x_i = -1, \quad v^i x_i = 0, \quad (3)$$

where ρ is the proper energy density for a cloud string with particles attached to them, λ is the string tension density, v^i is the four-velocity of the particles, and x^i is a unit space-like vector representing the direction of string. If the particle density of the configuration is denoted by ρ_p , then we have

$$\rho = \rho_p + \lambda. \quad (4)$$

The Einstein's field equations (in gravitational units $c = 1$, $G = 1$) read

$$R_i^j - \frac{1}{2} R g_i^j = -8\pi T_i^j, \quad (5)$$

where R_i^j is the Ricci tensor; $R = g^{ij} R_{ij}$ is the Ricci scalar. In a co-moving co-ordinate system, we have

$$v^i = (0, 0, 0, 1), \quad x^i = (0, 0, 1/C, 0). \quad (6)$$

The field equations (5) with Eq. (2) subsequently lead to the following system of equations:

$$\frac{B_{44}}{B} + \frac{C_{44}}{C} + \frac{B_4 C_4}{BC} = 8\pi \xi \theta, \quad (7)$$

$$\frac{A_{44}}{A} + \frac{C_{44}}{C} + \frac{A_4 C_4}{AC} = 8\pi \xi \theta, \quad (8)$$

$$\frac{A_{44}}{A} + \frac{B_{44}}{B} + \frac{A_4 B_4}{AB} - \frac{a^2}{A^2} = 8\pi(\lambda + \xi \theta), \quad (9)$$

$$\frac{A_4 B_4}{AB} + \frac{B_4 C_4}{BC} + \frac{C_4 A_4}{CA} - \frac{a^2}{A^2} = 8\pi \rho, \quad (10)$$

$$\frac{A_4}{A} - \frac{B_4}{B} = 0, \quad (11)$$

where the subscript 4 at the symbols A , B and C denotes ordinary differentiation with respect to t . The particle density ρ_p is given by

$$8\pi\rho_p = \frac{B_4C_4}{BC} + \frac{C_4A_4}{CA} - \frac{B_{44}}{B} - \frac{A_{44}}{A} + 8\pi K \quad (12)$$

in accordance with Eq. (4).

The velocity field v^i as specified by Eq. (6) is irrotational. The expansion (θ) and components of shear tensor (σ_{ij}) are given by

$$\theta = \frac{A_4}{A} + \frac{B_4}{B} + \frac{C_4}{C}, \quad (13)$$

$$\sigma^1{}_{-1} = \frac{1}{3} \left[\frac{2A_4}{A} - \frac{B_4}{B} - \frac{C_4}{C} \right], \quad (14)$$

$$\sigma^2{}_{-2} = \frac{1}{3} \left[\frac{2B_4}{B} - \frac{A_4}{A} - \frac{C_4}{C} \right], \quad (15)$$

$$\sigma^3{}_{-3} = \frac{1}{3} \left[\frac{2C_4}{C} - \frac{A_4}{A} - \frac{B_4}{B} \right], \quad (16)$$

$$\sigma^4{}_{-4} = 0. \quad (17)$$

Therefore

$$\sigma^2 = \frac{1}{2} [(\sigma^1{}_{-1})^2 + (\sigma^2{}_{-2})^2 + (\sigma^3{}_{-3})^2 + (\sigma^4{}_{-4})^2]$$

leads to

$$\sigma^2 = \frac{1}{3} \left[\frac{A_4^2}{A^2} + \frac{B_4^2}{B^2} + \frac{C_4^2}{C^2} - \frac{A_4B_4}{AB} - \frac{B_4C_4}{BC} - \frac{A_4C_4}{AC} \right]. \quad (18)$$

3 Solutions of the Field Equations

The field equations (7)-(11) are a system of five equations with six unknown parameters A , B , C , ρ , λ and ξ . One additional constraint relating these parameters is required to obtain explicit solutions of the system. Referring to Thorn [38], observations of the velocity-redshift relation for extragalactic sources suggest that Hubble expansion of the universe is isotropic today to within about 30 percent [39, 40]. Put more precisely, redshift studies place the limit

$$\frac{\sigma}{H} \leq 0.30$$

on the ratio of shear, σ , to Hubble constant H in the neighbourhood of our Galaxy. Following Bali and Jain [41] and Pradhan et al. [42, 43], we assume

that the expansion (θ) in the model is proportional to the shear (σ), which is physically plausible condition. This condition leads to

$$B = \ell C^m \quad (19)$$

where ℓ is proportionality constant and n is a constant. Equations (11) leads to

$$A = mB, \quad (20)$$

where m is an integrating constant. To obtain the determinate model of the universe, we assume that the coefficient of bulk viscosity is inversely proportional to expansion (θ). This condition leads to

$$\xi\theta = K, \quad (21)$$

where K is a proportionality constant.

By the use of Eqs. (19) and (21), Eq. (7) leads to

$$2C_{44} + \left(\frac{2n^2}{n+1}\right) \frac{C_4^2}{C} = \left(\frac{16\pi K}{n+1}\right) C. \quad (22)$$

Let

$$C_4 = f(C). \quad (23)$$

Hence Eq. (22) leads to

$$\frac{d}{dC} (f^2) + \left(\frac{2n^2}{n+1}\right) \frac{f^2}{C} = \left(\frac{16\pi K}{n+1}\right) C. \quad (24)$$

From Eq. (24), we have

$$f^2 C^{\frac{2n^2}{n+1}} = \left(\frac{8\pi K}{n^2+n+1}\right) C^{\frac{2n^2+2n+2}{n+1}} + L, \quad (25)$$

where L is an integrating constant. Eq. (25) leads to

$$f^2 = \left(\frac{8\pi K}{n^2+n+1}\right) C^2 + LC^{-(\frac{2n^2}{n+1})}. \quad (26)$$

From Eq. (26), we have

$$\frac{C^{\frac{n^2}{n+1}} dC}{\sqrt{C^2\left(\frac{n^2+n+1}{n+1}\right) + b^2}} = \sqrt{a} dt. \quad (27)$$

Integrating Eq. (27), we have

$$C = \left[\frac{\beta^2(\beta t + \gamma)^2 - b^2}{\beta^2} \right]^{\left(\frac{n+1}{n^2+n+1}\right)}, \quad (28)$$

where γ is an integrating constant and

$$\beta = \frac{(n^2 + n + 1)\sqrt{a}}{2(n + 1)},$$

$$a = \frac{8\pi K}{(n^2 + n + 1)},$$

$$b^2 = \frac{L}{a}.$$

Therefore, we have

$$B = \ell C^n = \ell \left[\frac{\beta^2(\beta t + \gamma)^2 - b^2}{\beta^2} \right]^{\left(\frac{n(n+1)}{n^2+n+1}\right)}, \quad (29)$$

$$A = mB = \ell m \left[\frac{\beta^2(\beta t + \gamma)^2 - b^2}{\beta^2} \right]^{\left(\frac{n(n+1)}{n^2+n+1}\right)}. \quad (30)$$

Hence the metric (1) reduces to the form

$$\begin{aligned} ds^2 = & -dt^2 + \left[\frac{\beta^2(\beta t + \gamma)^2 - b^2}{\beta^2} \right]^{\frac{2n(n+1)}{n^2+n+1}} [dX^2 + e^{-2\alpha X} dY^2] \\ & + \left[\frac{\beta^2(\beta t + \gamma)^2 - b^2}{\beta^2} \right]^{\frac{2(n+1)}{n^2+n+1}} dZ^2, \end{aligned} \quad (31)$$

where

$$\ell m x = X, \quad \ell y = Y, \quad z = Z.$$

Using the transformation

$$\beta(\beta t + \gamma) = b \cos(\beta \tau) \quad (32)$$

the metric (31) reduces to the form

$$\begin{aligned} ds^2 = & -\frac{b^2 \sin^2(\beta \tau)}{\beta^2} d\tau^2 + \left[\frac{b^4 \sin^4(\beta \tau)}{\beta^4} \right]^{\frac{n(n+1)}{n^2+n+1}} [dX^2 + e^{-\frac{2\alpha X}{\ell m}} dY^2] \\ & + \left[\frac{b^4 \sin^4(\beta \tau)}{\beta^4} \right]^{\frac{(n+1)}{n^2+n+1}} dZ^2. \end{aligned} \quad (33)$$

In absence of viscosity i.e. when $\beta \rightarrow 0$ then the metric (33) reduces to the new form

$$ds^2 = -b^2 \tau^2 d\tau^2 + (b^4 \tau^4)^{\frac{n(n+1)}{n^2+n+1}} [dX^2 + e^{-\frac{2\alpha X}{\ell m}} dY^2] + (b^4 \tau^4)^{\frac{(n+1)}{n^2+n+1}} dZ^2. \quad (34)$$

4 Some Physical and Geometrical Features of the Models

The rest energy density (ρ), the string tension density (λ), the particle density (ρ_p), the scalar of expansion (θ) and the shear (σ) for the model (31) are given by

$$8\pi\rho = \frac{4n(n+2)(n+1)^2}{(n^2+n+1)^2} \frac{\beta^2(\beta t + \gamma)^2}{\left[\frac{\beta^2(\beta t + \gamma)^2 - b^2}{\beta^2}\right]^2} - \frac{\alpha^2}{m^2\ell^2 \left[\frac{\beta^2(\beta t + \gamma)^2 - b^2}{\beta^2}\right]^{\frac{2n(n+1)}{n^2+n+1}}}, \quad (35)$$

$$8\pi\lambda = \frac{4n(n+1)\beta^2}{(n^2+n+1) \left[\frac{\beta^2(\beta t + \gamma)^2 - b^2}{\beta^2}\right]} + \frac{4n(n+1)\beta^2(\beta t + \gamma)^2(n^2+n-2)}{(n^2+n+1)^2 \left[\frac{\beta^2(\beta t + \gamma)^2 - b^2}{\beta^2}\right]^2} - \frac{\alpha^2}{m^2\ell^2 \left[\frac{\beta^2(\beta t + \gamma)^2 - b^2}{\beta^2}\right]^{\frac{2n(n+1)}{n^2+n+1}}} - \frac{\beta^2(n+1)^2}{(n^2+n+1)}, \quad (36)$$

$$8\pi\rho_p = \frac{8n(n+1)(n+2)\beta^2(\beta t + \gamma)^2}{(n^2+n+1)^2 \left[\frac{\beta^2(\beta t + \gamma)^2 - b^2}{\beta^2}\right]^2} - \frac{4n(n+1)\beta^2}{(n^2+n+1) \left[\frac{\beta^2(\beta t + \gamma)^2 - b^2}{\beta^2}\right]} + \frac{\beta^2(n+1)^2}{(n^2+n+1)}, \quad (37)$$

$$\theta = \frac{2(2n+1)(n+1)\beta(\beta t + \gamma)}{(n^2+n+1) \left[\frac{\beta^2(\beta t + \gamma)^2 - b^2}{\beta^2}\right]}, \quad (38)$$

$$\sigma = \frac{2(n^2-1)\beta(\beta t + \gamma)}{(n^2+n+1)\sqrt{3} \left[\frac{\beta^2(\beta t + \gamma)^2 - b^2}{\beta^2}\right]}. \quad (39)$$

The energy conditions $\rho \geq 0$ and $\rho_p \geq 0$ are satisfied in the presence of bulk viscosity for the model (31). The condition $\rho \geq 0$ leads to

$$-\frac{\alpha(n^2+n+1)}{2\ell m(n+1)\sqrt{n(n+2)}} \leq \frac{\beta(\beta t + \gamma)}{\left[\frac{\beta^2(\beta t + \gamma)^2 - b^2}{\beta^2}\right]^{\frac{1}{n^2+n+1}}} \leq \frac{\alpha(n^2+n+1)}{2\ell m(n+1)\sqrt{n(n+2)}}.$$

From Eq. (36), we observe that the string tension density $\lambda \geq 0$ provided

$$\frac{1}{\left[\frac{\beta^2(\beta t + \gamma)^2 - b^2}{\beta^2}\right]} + \frac{(\beta t + \gamma)^2(n^2+n-2)}{(n^2+n+1) \left[\frac{\beta^2(\beta t + \gamma)^2 - b^2}{\beta^2}\right]} \geq \frac{\alpha^2(n^2+n+1)}{4n(n+1)m^2\ell^2\beta^2 \left[\frac{\beta^2(\beta t + \gamma)^2 - b^2}{\beta^2}\right]^{\frac{2n(n+1)}{n^2+n+1}}} + \frac{(n+1)}{4n}.$$

The model in presence of bulk viscosity starts with a big bang at time $t = \frac{b}{\beta^2} - \frac{\gamma}{\beta}$. The expansion in the model decreases as time increases. The expansion in the model stops at $t = -\frac{\gamma}{\beta}$. The model (31) in general represents shearing and non-rotating universe. The role of bulk viscosity is to retard expansion in the model. We can see from the above discussion that the bulk viscosity plays a significant role in the evolution of the universe. Furthermore, since $\lim_{t \rightarrow \infty} \frac{\sigma}{\theta} \neq 0$, the model does not approach isotropy for large value of t . However the model isotropizes when $t = -\frac{\gamma}{\beta}$. There is a real physical singularity in the model (31) at $t = 0$.

Using the transformation

$$\beta(\beta t + \gamma) = b \cos(\beta \tau),$$

the rest energy density (ρ), the string tension density (λ), the particle density (ρ_p), the scalar of expansion (θ) and the shear (σ) for the model (33) are given by

$$8\pi\rho = \frac{4n(n+2)(n+1)^2 \cos^2(\beta\tau)}{b^2(n^2+n+1)^2 \left[\frac{\sin^4(\beta\tau)}{\beta^4} \right]} - \frac{\alpha^2}{m^2 \ell^2 \left[b^2 \frac{\sin^2(\beta\tau)}{\beta^2} \right]^{\frac{2n(n+1)}{n^2+n+1}}}, \quad (40)$$

$$8\pi\lambda = -\frac{4n(n+1)\beta^2}{(n^2+n+1) \left[b^2 \frac{\sin^2(\beta\tau)}{\beta^2} \right]} + \frac{4n(n+1)(n^2+n-2)b^2 \cos^2(\beta\tau)}{(n^2+n+1)^2 \left[b^2 \frac{\sin^2(\beta\tau)}{\beta^2} \right]^2} - \frac{\alpha^2}{m^2 \ell^2 \left[b^2 \frac{\sin^2(\beta\tau)}{\beta^2} \right]^{\frac{2n(n+1)}{n^2+n+1}}} - \frac{\beta^2(n+1)^2}{(n^2+n+1)}, \quad (41)$$

$$8\pi\rho_p = \frac{8n(n+1)(n+2)b^2 \cos^2(\beta\tau)}{(n^2+n+1)^2 \left[b^2 \frac{\sin^2(\beta\tau)}{\beta^2} \right]^2} + \frac{4n(n+1)\beta^2}{(n^2+n+1) \left[b^2 \frac{\sin^2(\beta\tau)}{\beta^2} \right]} + \frac{\beta^2(n+1)^2}{(n^2+n+1)}, \quad (42)$$

$$\theta = \frac{2(2n+1)(n+1) \cos(\beta\tau)}{N(n^2+n+1) \frac{\sin^2(\beta\tau)}{\beta^2}}, \quad (43)$$

$$\sigma = \frac{2(n^2-1)}{\sqrt{3}N(n^2+n+1) \frac{\sin^2(\beta\tau)}{\beta^2}}, \quad (44)$$

where $b = -N$, $N > 0$. In absence of bulk viscosity i.e. when $\beta \rightarrow 0$ then the physical and kinematic quantities for the model (34) are given by

$$8\pi\rho = \frac{4n(n+2)(n+1)^2}{b^2(n^2+n+1)^2} \frac{1}{\tau^4} - \frac{\alpha^2}{m^2 \ell^2} \frac{1}{[b^4 \tau^4]^{\frac{n(n+1)}{n^2+n+1}}}, \quad (45)$$

$$8\pi\lambda = \frac{4n(n+1)(n^2+n-2)}{b^2(n^2+n+1)^2 \tau^4} - \frac{\alpha^2}{m^2 \ell^2 [b^4 \tau^4]^{\frac{n(n+1)}{n^2+n+1}}}, \quad (46)$$

$$8\pi\rho_p = \frac{8n(n+1)(n+2)}{b^2(n^2+n+1)^2} \frac{1}{\tau^4}, \quad (47)$$

$$\theta = \frac{2(n^2 - 1)}{N(n^2 + n + 1)} \frac{1}{\tau^2}, \quad (48)$$

$$\sigma = \frac{2(n^2 - 1)}{N\sqrt{3}(n^2 + n + 1)} \frac{1}{\tau^2}. \quad (49)$$

In the absence of bulk viscosity, the energy conditions $\rho \geq 0$ and $\rho_p \geq 0$ are satisfied for the model (34). The condition $\rho \geq 0$ leads to

$$\tau^{\frac{4}{n^2+n+1}} \leq \frac{4n(n+1)^2(n+2)m^2\ell^2b^{2(n^2+n-1)}}{\alpha^2(n^2+n+1)^2}.$$

The string tension density $\lambda \geq 0$ if

$$\tau^{\frac{4}{n^2+n+1}} \leq \frac{4n(n+1)(n^2+n-2)m^2\ell^2b^{2(n^2+n-1)}}{\alpha^2(n^2+n+1)^2}.$$

The model (34) in the absence of bulk viscosity starts with a big bang at time $\tau = 0$ and the expansion in the model decreases as time increases. Since $\lim_{\tau \rightarrow \infty} \frac{\sigma}{\theta} \neq 0$, the model does not approach isotropy for large value of τ . However the model isotropizes for $n = -1$ and $n = 1$. There is a real physical singularity in the model (34) at $\tau = 0$.

Acknowledgments

We would like to thank the Inter-University Centre for Astronomy and Astrophysics (IUCAA), Pune, India for providing facility and support where this work was carried out. We also thank the anonymous referees for the fruitful comments.

References

- [1] Kibble, T. W. B. (1976). *J. Phys. A: Math. Gen.* **9**, 1387.
- [2] Zel'dovich, Ya. B., Kobzarev, I. Yu., and Okun, L. B. (1975). *Zh. Eksp. Teor. Fiz.* **67**, 3; (1975). *Sov. Phys.-JETP* **40**, 1.
- [3] Kibble, T. W. B. (1980). *Phys. Rep.* **67**, 183.
- [4] Everett, A. E. (1981). *Phys. Rev.* **24**, 858.
- [5] Vilenkin, A. (1981). *Phys. Rev. D* **24**, 2082.
- [6] Zel'dovich, Ya. B., (1980). *Mon. Not. R. Astron. Soc.* **192**, 663.
- [7] Letelier, P. S., (1979). *Phys. Rev. D* **20**, 1249.
- [8] Letelier, P. S., (1983). *Phys. Rev. D* **28**, 2414.

- [9] Stachel, J., (1980). *Phys. Rev.* **D 21**, 2171.
- [10] Banerjee, A., Sanyal, A. K. and Chakraborty, S., (1990). *Pramana-J. Phys.* **34**, 1
- [11] Chakraborty, S., (1991). *Ind. J. Pure Appl. Phys.* **29**, 31.
- [12] Tikekar, R. and Patel, L. K., (1992). *Gen. Rel. Grav.* **24**, 397; (1994). *Pramana-J. Phys.* **42**, 483.
- [13] Patel, L. K. and Maharaj, S. D., (1996). *Pramana-J. Phys.* **47**, 1.
- [14] Ram, S. and Singh, T. K., (1995). *Gen. Rel. Grav.* **27**, 1207.
- [15] Carminati, J. and McIntosh, C. B. G., (1980). *J. Phys. A: Math. Gen.* **13**, 953.
- [16] Krori, K. D., Chaudhury, T., Mahanta, C. R. and Mazumdar, A., (1990). *Gen. Rel. Grav.* **22**, 123.
- [17] Wang, X. X., (2003). *Chin. Phys. Lett.* **20**, 615.
- [18] Singh, G. P. and Singh, T., (1999). *Gen. Relativ. Gravit.* **31**, 371.
- [19] Lidsey, J. E., Wands, D. and Copeland, E. J., (2000). *Phys. Rep.* **337**, 343.
- [20] Baysal, H., Yavuz, I., Tarhan, I., Camci, U. and Yilmaz, I., (2001). *Turk. J. Phys.* **25**, 283.
- [21] Bali, R. and Dave, S., (2001). *Pramana-J. Phys.* **56**, 513.
- [22] Bali, R. and Upadhaya, R. D., (2003). *Astrophys. Space Sci.* **283**, 97.
- [23] Bali, R. and Dave, S., (2003). *Astrophys. Space Sci.* **288**, 503.
- [24] Bali, R. and Singh, D. K., (2005). *Astrophys. Space Sci.* **300**, 387.
- [25] Bali, R. and Anjali, (2006). *Astrophys. Space Sci.* **302**, 201.
- [26] Yavuz, İlhami, Yilmaz, İhsan and Baysal, Hüsnü, (2006). *Int. J. Mod. Phys.* **D 8**, 1365.
- [27] Yilmaz, I., (2006). *Gen. Relativ. Gravit.* **38**, 1397.
- [28] Selijak, U., Slosar, A. and McDonald, P., (2006). *J. Cosm. Astroparticle Phys.* **10**, 014.
- [29] Selijak, U. and Slosar, A., (2006). Preprint astro-ph/0604143.
- [30] Maharaj, S. D., Leach, P. G. L. and Govinder, K. S., (1995). *Pramana-J. Phys.* **44**, 511.

- [31] Yadav, M. K., Pradhan, A. and Singh, S. K., (2006). *Some Magnetized Bulk Viscous String Cosmological Models in General Relativity*, Preprint.
- [32] Wang, X. X., (2004). *Astrophys. Space Sci.* **293**, 933.
- [33] Wang, X. X., (2004). *Chin. Phys. Lett.* **21**, 1205.
- [34] Wang, X. X., (2005). *Chin. Phys. Lett.* **22**, 29.
- [35] Wang, X. X., (2006). *Chin. Phys. Lett.* **23**, 1702.
- [36] Yadav, M. K., Rai, Anju, and Pradhan, A., (2006). *Int. J. Theor. Phys.*, Accepted. Preprint gr-qc/ 0611032.
- [37] Landau, L. D. and Lifshitz, E. M., (1963). *Fluid Mech.* **6**, 505.
- [38] Thorne, K. S., (1967). *Astrophys. Space Sci.* **148**, 51.
- [39] Kantowski, R. and Sachs, R. K., (1966). *J. Math. Phys.* **7**, 433.
- [40] Kristian, J. and Sachs, R. K., (1966). *Astrophys. J.* **143**, 379.
- [41] Bali R. and Jain, V. C., (1999). *Astrophys. Space Sci.* **262**, 145.
- [42] Pradhan, A. and Yadav, L. and Yadav, A. K., (2003). *Czech. J. Phys.* **54**, 487.
- [43] Pradhan, A. and Singh, S. K., (2004). *Int. J. Mod. Phys.* **D13**, 503.